arXiv:alg-geom/9711010v1 10 Nov 1997

Constructing Curves over Finite Fields with Many Points by Solving Linear Equations

Gerard van der Geer & Marcel van der Vlugt

The purpose of this note is to exhibit an elementary method to construct explicitly curves over finite fields with many points. Despite its elementary character the method is very efficient and can be regarded as a partial substitute for the use of class field theory. Many of the results on the existence of curves with a large number of points obtained from class field theory or Drinfeld modules can thus be reproduced with *explicit* curves and many new examples can easily be obtained. Curves with many points find applications in coding theory and the theory of low-discrepancy sequences and here explicitness is often essential.

The Method

Let $k = \mathbb{F}_q$ be a finite field of cardinality $q = p^m$ with p prime. We consider fibre products C_F over a base curve C of Artin-Schreier extensions $C_f \to C$ defined by function field extensions $k(C_f) = k(C)(z)$, where z satisfies an equation $z^p - z = f$ and f runs through a basis of functions $f \in F$ with F a suitable $\mathbb{F}_p$-vectorspace of $k(C)$. In order to get a curve C_F with many points we shall start with a curve C over k with many rational points. We select a (large) set of distinct rational points $\mathcal{P}$ on C and an effective divisor D defined over k whose support is disjoint from $\mathcal{P}$. We consider a $\mathbb{F}_p$-vector space

$$F \subseteq L(D) = \{g \in k(C) : (g) + D \geq 0\} \cup \{0\}$$

of functions f satisfying

$$\begin{aligned} &\text{i)} \quad F \cap \{g^p - g : g \in k(C)\} = \{0\}, \\ &\text{ii)} \quad \mathrm{Tr}_{q/p}(f(P)) = 0 \quad \text{for all } P \in \mathcal{P}, \end{aligned}$$

where $\mathrm{Tr}_{q/p}$ is the trace map from $\mathbb{F}_q$ to $\mathbb{F}_p$. The fibre product C_F is a covering of C in which all the rational points $P \in \mathcal{P}$ are completely split by ii), so that we may expect many points. The genus of C_F is determined by the genera of the building blocks C_f with $f \in F - \{0\}$ and these follow from the Hurwitz-Zeuthen formula. A sensible choice of D must be made in order to keep the genus of C_F relatively small. Linear algebra gives estimates for the dimensions of subspaces F satisfying i) and ii).

Covers of Curves

Let C be a smooth complete irreducible curve defined over k and suppose that $f_1, \ldots, f_r$ is a basis of a space of functions F satisfying condition i). Each curve C_f defined by the equation $z_i^p - z_i = f_i$ is an irreducible cover of C; up to C-isomorphism the fibre product

$$C_F = C_{f_1} \times_C \times \ldots \times_C C_{f_r}$$

does not depend on the chosen basis. The Jacobian $J(C_F)$ of C_F decomposes up to isogeny as product

$$J(C) \times \prod_{f \in \mathbb{P}(F)} P_f,$$

where P_f is the Prym variety of $C_f \to C$ (i.e. the connected component of the kernel of the norm map $\mathrm{Nm} : J(C_f) \to J(C)$). The sum is over a complete set of representatives $f \in F$ modulo the equivalence relation $f \sim c\, f$ with $c \in \mathbb{F}_p^*$ (which we write as $f \in \mathbb{P}(F)$).

If we denote the trace of Frobenius of C (resp. C_f, C_F) by τ (resp. τ_f, τ_F) then we find the relation

$$\tau_F = \tau + \sum_{f \in \mathbb{P}(F)} (\tau_f - \tau)$$

from which we can compute the number of points of C_F. The genus of C_F is given similarly by

$$g(C_F) = g(C) + \sum_{f \in \mathbb{P}(F)} (g(C_f) - g(C)).$$

Curves arising from Spaces of Solutions

To guarantee that we have sufficiently many functions we use Riemann-Roch. Let $\mathcal{P} = \{P_1, \ldots, P_n\}$ be the set of $k = \mathbb{F}_q$-rational points on C. For an effective divisor $D = \sum n_Q Q$ defined over $\mathbb{F}_q$ we denote by $[D/p]$ the divisor $\sum [n_Q/p] Q$ with $[x]$ the greatest integer function. We set

$$\delta = \#(\mathrm{Supp}(D) \cap \mathcal{P}).$$

Proposition 1. *For a curve C and a divisor D as above there exists for every integer r with*

$$1 \leq r \leq \dim_{\mathbb{F}_p}(L(D)) - \dim_{\mathbb{F}_p}(L([D/p]) + 1 - n + \delta$$

a $\mathbb{F}_p$-subspace $F_r \subset L(D)$ of dimension r such that for $f \in F_r - \{0\}$ the curve C_f satisfies

$$\#C_f(\mathbb{F}_q) = p(n - \delta) + \epsilon_f,$$

where $\epsilon_f \geq 0$ is the number of rational points of C_f lying over points in $\mathrm{Supp}(D)$.

Proof. Let $\bar{\mathbb{F}}$ denote an algebraic closure of $\mathbb{F}_q$. From [V] it follows that the $\mathbb{F}_p$-vector space

$$V = \{f \in L(D) \colon f = g^p - g \text{ for some } g \in \bar{\mathbb{F}}(C)\}$$

has dimension $\dim_{\mathbb{F}_p}(L([D/p]))$. Choose now a linear $\mathbb{F}_p$-space $W \subset L(D)$ with $W \cap V = \{0\}$ and with dimension $\dim_{\mathbb{F}_p}(L(D)) - \dim_{\mathbb{F}_p}(L([D/p]))$. Let $c \in \mathbb{F}_q$ with $\mathrm{Tr}_{q/p}(c) \neq 0$, set

$$\tilde{W} = W \oplus \mathbb{F}_p \cdot c$$

and solve in $\tilde{W}$ the linear equations

$$\mathrm{Tr}_{q/p}(f(P_i)) = 0 \quad \text{for all } P_i \notin \mathrm{Supp}(D). \tag{1}$$

The solution space of this homogeneous system of linear equations over $\mathbb{F}_p$ has dimension $\geq \dim \tilde{W} - (n - \delta)$. The equations (1) express the condition that all the points P_i which do not belong to $\mathrm{Supp}(D)$ are completely split. This gives the expression for $\#C_f(\mathbb{F}_q)$. □

We now form the fibre product C_F associated to a space of solutions F of (1) and we obtain:

Corollary. *With the notations as in Proposition 1 there exists for every r with*

$$1 \leq r \leq \dim_{\mathbb{F}_p}(L(D)) - \dim_{\mathbb{F}_p}(L([D/p])) + 1 - n + \delta$$

a r-dimensional subspace $F_r \subset L(D)$ such that

$$\#C_{F_r}(\mathbb{F}_q) = p^r(n - \delta) + \delta + \sum_{f \in \mathbb{P}(F_r)} (\epsilon_f - \delta).$$

Remark. For $f \in F_r, f \neq 0$ the Hurwitz-Zeuthen formula implies:

$$2g(C_f) - 2 = p(2g(C) - 2) + (p - 1) \sum_{Q} ((-v_Q^*(f) + 1) \deg(Q)),$$

where the summation is over those Q in $\mathrm{Supp}(D)$ with $-v_Q^*(f) > 0$. Here $v_Q^*(f)$ denotes the Artin-Schreier reduced order at the pole of f in Q.

Examples

Examples can now be obtained in a straightforward way. By considering the special cases

$$\begin{aligned} D &= 2Q && \text{with } Q \text{ a point of degree } d \text{ on } C, \\ D &= kP && \text{with } k \in \mathbb{Z}_{>0} \text{ and } P \text{ a rational point of } C, \\ D &= 2P + 2Q && \text{with } P \text{ and } Q \text{ as above,} \end{aligned}$$

one can reproduce, but in *explicit form*, many of the examples given in the papers of Niederreiter and Xing (see References). There the curves were implicitly given by subfields of narrow ray class fields determined by Drinfeld modules of rank 1 over a curve. However, if the base curve has genus ≥ 1 it is rather difficult to find explicit forms of the curves. The same holds for examples constructed by class field theory in [L], [Sch] and [S1,2].

Example 1. In [S3] Serre constructed curves C over $\mathbb{F}_2$ with invariants $(g(C), \#C(\mathbb{F}_2)) = (10, 13), (11, 14)$ and $(13, 15)$ as covers of the elliptic curve E defined by $y^2+y = x^3+x$ using the group structure of $E(\mathbb{F}_2)$. We show how these examples can be recovered with our method. The rational points of E are P_∞, $P_1 = (0, 0)$, $P_2 = (1, 0)$, $P_3 = (1, 1)$ and $P_4 = (0, 1)$. Consider the space $\tilde{W}$ for $D = 3P_\infty + P_1 + P_3$ with $\mathbb{F}_2$-basis

$$1, x, y, x/(x + y), xy/(x + y).$$

We now require in $\tilde{W}$ that P_2 and P_4 are completely split. The solution space of (1) has basis $f_1 = x + x/(x + y)$, $f_2 = 1 + x + y$ and $f_3 = xy/(x + y)$. The 2-dimensional $\mathbb{F}_2$-subspace $F = \langle f_1, f_2 + f_3\rangle$ yields a curve of genus 10 with 13 points.

With $D = 3P_\infty + 3P_4$ we can take $\tilde{W} = \langle 1, y/x, x, y, y/(x(x + y + 1)\rangle$. Then $F = \langle 1 + y/x + x + y, (xy + y^2)/x(x + y + 1)\rangle$ yields the example with $(g, \#) = (11, 14)$.

With $D = 7P_\infty + P_4$ and splitting $\{P_1, P_2, P_3\}$ we find a 3-dimensional space $H = \langle 1 + x + y + y/x, y + xy, y + x^2y\rangle$. The subspace $F = \langle 1 + x + y + y/x, xy + x^2y\rangle$ produces a curve C_F with $(g(C_F), \#C_F(\mathbb{F}_2)) = (13, 15)$. Other 2-dimensional spaces produce curves with $(g, \#) = (11, 13), (12, 13), (13, 13), (14, 13), (14, 14)$ and $(14, 15)$. The curve C_H has $(g(C_H), \#C_H(\mathbb{F}_2)) = (29, 25)$.

Example 2 i). Consider the elliptic curve C defined by $y^3 - y = x^2 - 1$ over $\mathbb{F}_3$ with $\#C(\mathbb{F}_3) = 7$. Letting $P_\infty = (0 : 1 : 0)$, $Q = (1, 0)$ and $D = 8P_\infty + 2Q$ we can take $\tilde{W}$ with $\mathbb{F}_3$-basis

$$1, (y + 2)/(2x + y + 1), (x + y + 1)/y, y, x + 2, y^2, (x + 2)y, (x + 2)y^2, (x + 2)^2y.$$

We solve in $\tilde{W}$ for the remaining 5 rational points P_i $(i = 1, \ldots, 5)$ of C the equations

$$\mathrm{Tr}_{3/3}(f(P_i)) = f(P_i) = 0 \qquad i = 1, \ldots, 5$$

and we get a 4-dimensional solution space F generated by

$$f_1 = \frac{y + 2}{2x + y + 1} + y + x + y^2, \quad f_2 = \frac{y + 2}{2x + y + 1} + \frac{x + y + 1}{y} + x + 2xy,$$
$$f_3 = 2 + \frac{y + 2}{2x + y + 1} + y + 2y^2+xy^2, \quad \text{and} \quad f_4 = 2y + x^2y$$

with the following curves

$f_i =$	$g(C_{f_i})$	$\#C_{f_i}(\mathbb{F}_3)$
f_1	9	17
f_2	10	17
f_3	12	17
f_4	10	19

Analyzing the subspaces of F we obtain the following table, in which under $N_q(g)$ we list the best curve so far and the Oesterlé upper bound (cf. [G-V]).

F	$g(C_F)$	$\#C_F(\mathbb{F}_3))$	$N_q(g)$
$\langle f_1, f_2\rangle$	35	47	$[38-51]$
$\langle f_1+2f_3, f_4\rangle$	36	46	$[36-52]$
$\langle f_1, f_3\rangle$	39	46	$[42-56]$
$\langle f_1, f_2, f_3\rangle$	128	136	≤ 149

Example 2 ii). Consider again the elliptic curve C given by $y^3 - y = x^2 - 1$ over $\mathbb{F}_3$ with $\#C(\mathbb{F}_3) = 7$. Let D be the divisor $3Q$ with Q a point of degree 4 on C given by the ideal (x^2+xy+1, x^4+x^3+2). This is the Frobenius orbit of $(\alpha, \alpha^3+\alpha^9)$. Requiring that the 7 rational points are completely split leads to a 2-dimensional space F with basis

$$1 + f_2 + 2f_3 + 2f_1^2 + f_2^2 + f_3^2 \qquad \text{and} \qquad 2 + f_1^2 + f_2^2 + f_1 f_2 f_3,$$

where $f_1 = (2x+y)/(x^2+xy+1)$, $f_2 = (2x^2+1)/(x^2+xy+1)$, $f_3 = (2x^2+y^2)/(x^2+xy+1)$. Then C_F is a curve of genus 49 and $\#C_F(\mathbb{F}_3) = 63$. (Best result so far $N_q(g) \in [49-67]$.)

Example 3. Let $\mathbb{F}_4 = \mathbb{F}_2(\alpha)$ and let C be the elliptic curve defined by $y^2 + y = x^3$ with $\#C(\mathbb{F}_4) = 9$. We use $P_\infty = (0:1:0)$ and set $D = 11P_\infty$. Elementary calculations give $\dim_{\mathbb{F}_2}(\tilde{W}) = 13$ and we may take as a $\mathbb{F}_2$-basis:

$$\tilde{W} = \langle \alpha, x, \alpha x, y, \alpha y, xy, \alpha xy, x^2 y, \alpha x^2 y, y^3, \alpha y^3, xy^3, \alpha xy^3\rangle.$$

Splitting the 8 rational points different from P_∞ yields 8 equations in 12 variables with a 5-dimensional solution space. We list properties of the curves C_{f_i} corresponding to a basis f_i $(i = 1, \ldots, 5)$ of the solution space in the following table:

$f_i =$	$g(C_{f_i})$	$\#C_{f_i}(\mathbb{F}_4)$
$x + xy + x^2y$	5	17
$\alpha x + \alpha xy + \alpha^2 x^2 y$	5	17
y^3	6	17
$x + xy^3$	7	17
$\alpha x + \alpha xy^3$	7	17

The corresponding fibre products yield the following harvest:

F	$g(C_F)$	$\#C_F(\mathbb{F}_4))$	$N_q(g)$	comments
$\langle f_1, f_2\rangle$	13	33	33	best possible
$\langle f_1, f_3\rangle$	15	33	≤ 37	cf. Ex. 4.14 in [N-X1]
$\langle f_1, f_2, f_3\rangle$	33	65	≤ 66	cf. [L]
$\langle f_1, f_2, f_4\rangle$	37	65	≤ 72	cf. [N-X2]
$\langle f_1, f_3, f_4\rangle$	39	65	≤ 75	new

By employing this method systematically one can expect a substantial number of examples improving the tables (cf. [G-V]). We refer to work in progress by V. Shabat.

References

[G-V] G. van der Geer, M. van der Vlugt : Tables for the function $N_q(g)$. Regularly updated tables at: `http://www.wins.uva.nl/ ~geer` .

[L] K. Lauter: Ray class field constructions of curves over finite fields with many rational points. In: *Algorithmic Number Theory* (Talence 1996), H. Cohen Ed., Lecture Notes in Computer Science 1122, Springer, Berlin, 1996, p. 187-195.

[N-X1] H. Niederreiter, C. P. Xing: Cyclotomic function fields, Hilbert class fields and global function fields with many rational places. *Acta Arithm.* **79** (1997), p. 59-76.

[N-X2] H. Niederreiter, C. P. Xing: Drinfeld modules of rank 1 and algebraic curves with many rational points II. *Acta Arithm.* **81** (1997), p. 81-100.

[Sch] R. Schoof: Algebraic curves and coding theory. UTM 336, Univ. of Trento, 1990.

[S1] J-P. Serre : Sur le nombre de points rationnels d'une courbe algébrique sur un corps fini. *C.R. Acad. Sci. Paris* **296**, Série I (1983), p. 397-402. (= Oeuvres III, No. 128, p. 658-663).

[S2] J-P. Serre: Rational points on curves over finite fields. Notes of lectures at Harvard University 1985.

[S3] J-P. Serre: Letter to G. van der Geer, 1 September 1997.

[V] M. van der Vlugt: A new upper bound for the dimension of trace codes. *Bull. London Math. Soc.* **23** (1991), p. 395-400.

[X-N] C. P. Xing, H. Niederreiter: Modules de Drinfeld et courbes algébriques ayant beaucoup de points rationnels. *C.R. Acad. Sci. Paris* **322**, Série I (1996), p. 651-654.

G. van der Geer
Faculteit Wiskunde en Informatica
Universiteit van Amsterdam
Plantage Muidergracht 24
1018 TV Amsterdam
The Netherlands
`geer@wins.uva.nl`

M. van der Vlugt
Mathematisch Instituut
Rijksuniversiteit te Leiden
Niels Bohrweg 1
2333 CA Leiden
The Netherlands
`vlugt@wi.leidenuniv.nl`